

# TEGOSOFT<sup>®</sup> AC

A vegetable-based emollient with light skin feel

- Clear, low-viscous emollient that provides a nonoily skin feel and quick absorption into the skin
- Helps in formulating natural cosmetic products with a more light sensory profile
- Fully based on renewable sources
- Produced by an enzymatic process that leads to savings of >60% on energy consumption and  $CO_2$  emission
- NaTrue and COSMOS certified, approved by Ecocert

**Personal Care** 

#### INCI name

Isoamyl Cocoate

Chemical and physic (not part of specifica	

Form	Liquid

## Further product information (not part of specifications)

Density (g/cm <sup>3</sup> )	approx. 0.854
Viscosity at 25 °C according to Höppler (mPas)	approx. 6
Surface tension at 25 °C according to ring method (mN/m)	approx. 28
Spreadability	medium spreading
Polarity	medium polarity
Pour point (°C)	-3 ℃

#### Properties

- TEGOSOFT® AC is a low-viscous emollient. It can be easily distributed on the skin and it shows a quick absorption behavior without leaving a waxy film on the skin.
- It imparts a light, non-oily skin feel in cosmetic emulsions.
- It is miscible with all common cosmetic oils and can easily be used in O/W and W/O formulations.
- TEGOSOFT® AC is an emollient ester of isoamyl alcohol and fatty acid of 100% vegetable origins which are sugar beet, palm and coconut.
- It is produced by the eco-efficient Evonik enzymatic process which yields a high-purity product.
- A lifecycle assessment shows general savings in e. g. energy consumption and CO<sub>2</sub> emission comparing the enzymatic with the conventional esterification process.<sup>[1]</sup>

[1] O. Thum, K. M. Oxenboll SÖFW, 2008, 134, 44-47

#### Formulation hints

Slight hydrolyzation of isoamyl esters can occur at pH below 7. This has no effect on the stability of the emollient or formulation and in perfumed formulations this effect is negligible. In unperfumed formulation a odor of isoamyl alcohol could be perceived. This has be taken into account e.g. formulating with natural preservatives.

For the combination of isoamyl esters and UV filters the patent EP 2421614 has to be considered.

#### Applications

- Natural cosmetic products (see formulation hints)
- Facial and body care emulsions, gels and serums
- Baby Care
- Sprayable emulsions
- Antiperspirants/Deodorants
- Color cosmetics

#### Suggested usage concentration

1 - 10% TEGOSOFT® AC, up to 30% for make-up

#### Packaging

720 kg pallet (4 x 180 kg drum)

#### Storage

Store at max. 25 °C

#### Hazardous goods classification

Information concerning

- classification and labeling according to regulations for transport and for dangerous substances
- protective measures for storage and handling
- measures in accidents and fires
- toxicity and ecological effects

is given in our material safety data sheet.

### **Guideline Formulations**

O/W Serum	
(MK 10/10-4)	
Phase A	
TEGO <sup>®</sup> Care 165	3.00%
(Glyceryl Stearate; PEG-100 Stearate)	
TEGO <sup>®</sup> Alkanol 18 (Stearyl Alcohol)	0.50%
TEGOSOFT <sup>®</sup> AC (Isoamyl Cocoate)	5.50%
TEGOSOFT <sup>®</sup> CT	5.00%
(Caprylic/Capric Triglyceride)	
TEGOSOFT <sup>®</sup> CR (Cetyl Ricinoleate)	1.00%
Phase B	
Water	76.125%
Glycerin	3.00%
TEGO® Pep 4–17 (Tetrapeptide–21; Glycerin; Butylene Glycol; Aqua)	2.50%
TEGO <sup>®</sup> Pep 4–Even	2.50%
(Tetrapeptide-30, Glycerin)	
TEGO <sup>®</sup> Carbomer 141 G (Carbomer)	0.075%
Phase C	
Sodium Hydroxide (10% in water)	q.s.
Phase D	
Dipropylene Glycol; Methylparaban; Ethylparaben; Aqua; Methylisothiazolinone (Microcare MEM, Thor)	0.80%
Processing:	
<ol> <li>Heat phase B to approx. 75 °C and dispers Carbomer 141 G.</li> <li>Heat phase A separately to approx. 75 °C.</li> <li>Add phase A to phase B with stirring.<sup>1)</sup></li> <li>Homogenize.</li> <li>Cool with gentle stirring.</li> <li>Add phases C and D below 40 °C.</li> </ol>	e TEGO®
1)Important:	

<sup>1)</sup>Important: If phase A has to be charged into the vessel first, phase B must be added <u>without stirring</u>.

Age Defense BB Cream SPF 15 (MK 2/12-8)	
Phase A	
TEGIN <sup>®</sup> 4100 Pellets (Glyceryl Stearate)	1.00
Stearic Acid	1.00
TEGO <sup>®</sup> Alkanol 18 (Stearyl Alcohol)	3.00
TEGOSOFT® DEC	6.90
(Diethylhexyl Carbonate)	2.00
HyaCare <sup>®</sup> Filler CL (Aqua; Ethylhexyl Stearate; Sodium Hyaluronate Crosspolymer; Polyglyceryl-4 Diisostearate/Polyhydroxystearate/Sebacate; Sodium Isostearate)	2.00
Phytosphingosine (Phytosphhingosine)	0.10
Ethylhexyl Methoxycinnamate	5.00
Diethylamino Hydroxybenzoyl Hexyl	3.00
Benzoate	
Phase B	
Titanium Dioxide, Alumina, Triethoxycaprylylsilane (Hombitan AC 360, Sachtleben)	3.00
Talc (Micro Talc IT Extra-AW, Mondo Minerals)	2.00
Unipure Yellow LC 182 (Sensient)	0.36
Unipure Red LC 381 (Sensient)	0.12
Unipure Black LC 989 (Sensient)	0.08
TEGOSOFT <sup>®</sup> AC (Isoamyl Cocoate)	4.44
TEGOSOFT® XC (Phenoxyethyl Caprylate)	4.00
TEGOLON® ECO 10-10 (Nylon-10/10)	3.00
Phase C	
Water	51.10
Glycerin	3.00
TEGO° Care CG 90 (Cetearyl Glucoside)	1.00
HyaCare <sup>®</sup> 50 (Hydrolyzed Hyaluronic Acid)	0.10
TEGO® Pep 4–17 (Tetrapeptide–21; Glycerin; Butylene Glycol; Aqua)	2.00
Phase D	
Alcohol	3.00
Phase E	
Dipropylene Glycol; Methylparaban; Ethylparaben; Aqua; Methylisothiazolinone (Microcare MEM, Thor)	0.80
Processing:	
<ol> <li>Disperse the pigments/powders homogene phase B.</li> <li>Heat phase A and C separately to approx.</li> <li>Add phase A to phase C with stirring.<sup>1)</sup></li> <li>Homogenize.</li> <li>Add phase B to phase A/C at approx. 60 °C</li> <li>Homogenize for a short time.</li> <li>Cool with gentle stirring to approx. 40 °C a phase D and E.</li> </ol>	75 ℃. 2.
<sup>1)</sup> Important: If phase A has to be charged into the vessel fi B must be added <u>without stirring</u> .	rst, phase

Mattifying Compact Cream Make-up (AL 1/13-13)	
Phase A	
TEGOSOFT <sup>®</sup> AC (Isoamyl Cocoate)	31.70%
TEGOSOFT <sup>®</sup> XC (Phenoxyethyl Caprylate)	13.00%
TEGOSOFT <sup>®</sup> OP (Ethylhexyl Palmitate)	8.00%
TEGOSOFT <sup>®</sup> CR (Cetyl Ricinoleate)	5.00%
ABIL® Wax 9840 (Cetyl Dimethicone)	1.00%
Helianthus Annuus (Sunflower Seed) Wax	4.00%
Ozokerite	2.00%
Phase B	
Titanium Dioxide (Kronos 1171, Kronos)	12.00 %
Yellow Iron Oxide (Gelb 10 E, Sicovit)	1.00%
Brown Iron Oxide (Braun 70 E, Sicovit)	0.60%
Red Iron Oxide (Rot 30 E, Sicovit)	0.30%
Black Iron Oxide (Schwarz 80 E, Sicovit)	0.20%
TEGOLON <sup>®</sup> ECO 10-10 (Nylon-10/10)	20.00%
Phase C	
Glyceryl Caprylate (Dermosoft GMCY, Dr. Straetmans)	1.00%
Perfume	0.20%
Processing:	
<ol> <li>Heat phase A to 82 °C.</li> <li>Premix and grind phase B.</li> <li>Add phase B to phase A and stir gently at 80 °C</li> </ol>	

- Add phase b to phase A and still gently at ob e several minutes.
   Cool to 75 °C with gentle stirring.
   Add phase C with gentle stirring.
   Pour mixture into moulds and cool down to room to room

- temperature.

Skin Firming Body Lotion (MM 221/2)	
Phase A	
ISOLAN® GPS (Polyglyceryl-4 Diisostearate/Polyhydroxystearate/ Sebacate)	2.5%
TEGOSOFT <sup>®</sup> CT (Caprylic/Capric Triglyceride)	7.5%
TEGOSOFT <sup>®</sup> AC (Isoamyl Cocoate)	6.0%
Simmondsia Chinensis (Jojoba) Seed Oil	5.0%
Beeswax	0.6%
Hydrogenated Castor Oil	0.4%
Phase B	
Glycerin	3.0%
Magnesium Sulfate Heptahydrate	1.0%
Water	71.0%
Phase C	
TEGO <sup>®</sup> Stemlastin (Algae Extract)	3.0%
Phase Z	
Preservative, perfume	q.s.
Processing:	
<ol> <li>Heat phase A to approx. 80 °C.</li> <li>Add phase B (80 °C or room temperature) slip</li> </ol>	owly

- while stirring.
   Homogenize for a short time.
   Cool with gentle stirring and add phase C below 40 °C.
- 5. Homogenize again below 30 °C.

lcy and Smooth After Sun Gel (SG 44/11-7)	
Phase A	
TEGOSOFT® SH (Stearyl Heptaonoate)	5.00%
TEGOSOFT® AC (Isoamyl Cocoate)	5.00%
TEGO <sup>®</sup> Carbomer 341 ER (Acrylates/C10- 30 Alkyl Acrylate Crosspolymer)	0.30%
TEGO <sup>®</sup> Carbomer 140 (Carbomer)	0.30%
TEGOLON <sup>®</sup> ECO 10-10 (Nylon-10/10)	0.30%
Phase B	
Water	81.90%
Glycerin	2.00%
HyaCare <sup>®</sup> (Sodium Hyaluronate)	0.10%
TEGO <sup>®</sup> Cosmo C 100 (Creatine)	0.50%
Phase C	
Alcohol	2.00%
Phase D	
Sodium Hydroxide (10% in water)	1.80%
Phase E	
Dipropylene Glycol, Methylparaben, Ethylparaben, Aqua, Methyliso- thiazolinone (Microcare MEM, Thor)	0.80%
Phase Z	
Perfume	q.s.
<ol> <li>Processing:</li> <li>1. Disperse the powders in phase A.</li> <li>2. Mix ingredients of phase B separately.</li> <li>3. Combine both phases and homogenize.</li> <li>4. Add phase C and stir well.</li> </ol>	

4. Add phase C and stir well.

5. Add phase D and adjust the pH value to approx. 7.0.

6. Add phase E while stirring.

H 11/13

This information and all further technical advice is based on our present knowledge and experience. However, it implies no liability or other legal responsibility on our part, including with regard to existing third party intellectual property rights, especially patent rights. In particular, no warranty, whether express or implied, or guarantee of product properties in the legal sense is intended or implied. We reserve the right to make any changes according to technological progress or further developments. The customer is not released from the obligation to conduct careful inspection and testing of incoming goods. Performance of the product described herein should be verified by testing, which should be carried out only by qualified experts in the sole responsibility of a customer. Reference to trade names used by other companies is neither a recommendation, nor does it imply that similar products could not be used. (Status: April, 2008)

**Evonik Industries AG** Goldschmidtstraße 100 45127 Essen, Germany P.O. BOX 45116 Essen PHONE +49 201 173-2854 FAX +49 201 173-1828 personal-care@evonik.com www.evonik.com/personal-care

